

Composition and structure of the meteoritic macromolecule: A source of complex organics on the early Earth

Conel M. O'D. Alexander
*Dept. Terrestrial Magnetism
Carnegie Institution of Washington
5241 Broad Branch Road, NW
Washington DC 20015
USA
alexande@dtm.ciw.edu*

George D. Cody
*Geophysical Laboratory
Carnegie Institution of Washington
5251 Broad Branch Road, NW
Washington DC 20015
USA*

Marilyn Fogel
*Geophysical Laboratory
Carnegie Institution of Washington
5251 Broad Branch Road, NW
Washington DC 20015
USA*

Simon Kelemen
*ExxonMobil Research & Engineering Co.
Room LF-350
Annandale, NJ 08801
USA*

David Kilcoyne
*Advanced Light Source
Lawrence Berkeley Laboratory
1 Cyclotron Road
Berkeley, CA 94720
USA*

Tohru Araki
*Advanced Light Source
Lawrence Berkeley Laboratory
1 Cyclotron Road
Berkeley, CA 94720
USA*

Chondritic meteorites and interplanetary dust particles (IDPs) were a source of complex organic material on the early Earth. L enantiomer excesses in meteoritic amino acids has led to suggestions that this exogenous material played a role in the origin of life.

The majority of the organic material in chondrites is present as an insoluble, macromolecular material (IMM). The IMM's isotopic composition and IR spectrum suggests a direct link to refractory organic material in the diffuse interstellar medium. If the IMM played a role in the origin of life on Earth, its ubiquity in the Galaxy means it could be important to the development of life in other solar systems. Its composition also resembles that of the refractory organic matter in comet Halley dust.

IDPs may contain largely pristine IMM. In chondrites, aqueous alteration and thermal metamorphism modified the IMM to varying degrees. Many of the soluble organic compounds in chondrites may be alteration products of the IMM. Terrestrial weathering today rapidly breaks down the IMM. Weathering of the IMM on the early Earth may have been a more important source of complex soluble organics than those delivered directly by meteorites.

To understand the origin, chemistry and structure of the IMM, how it weathers and to quantify microanalytic techniques for STARDUST sample analysis, we have been conducting a multidisciplinary (NMR, XANES, XPS, EA and GC-MS) study of IMM. We will present our results for 32 chondrites, which reveal a complex functional group chemistry and behavior during aqueous alteration and metamorphism.